



A Survey for some Special Curves in Isotropic Space \mathbf{I}_3^1

Alper Osman Ogrenmis^{1*}, Mihriban Kulahci¹ and Mehmet Bektas¹

¹Department of Mathematics, Faculty of Science, Firat University, Elazig / TURKEY.

Authors' contributions

This work was carried out in collaboration between all authors. Author AOO designed the study and wrote the first draft of the manuscript. Authors MK and MB managed the analyses of the study. Author MK managed the literature searches. All authors read and approved the final manuscript.

Research Article

Received 22nd March 2013
Accepted 27th May 2013
Published 3^d June 2013

ABSTRACT

In this paper, curves of AW(k)-type in isotropic space \mathbf{I}_3^1 are defined. Using Frenet frames in isotropic space \mathbf{I}_3^1 , curvature conditions of AW(k)-type curves are given. In addition, new characterizations of Bertrand and Mannheim curves are obtained.

Keywords: Isotropic space; frenet frame; bertrand curves; mannheim curves; curvature; torsion.

1. INTRODUCTION

The assumption that our universe is homogeneous and isotropic means that its evolution can be represented as a time-ordered sequence of three-dimensional space-like hypersurfaces, each of which is homogeneous and isotropic. These hypersurfaces are the natural choice for surfaces of constants time.

Homogeneity means that the physical conditions are the same at every point of any given hypersurface. Isotropy means that the physical conditions are identical in all directions when

*Corresponding author: Email: ogrenmisalper@gmail.com;

viewed from a given point on the hypersurface. Isotropy at every point automatically enforces homogeneity. However, homogeneity does not necessarily imply isotropy.

Homogeneous and isotropic spaces have the largest possible symmetry group; in three dimensions there are three independent translations and three rotations. These symmetries strongly restrict the admissible geometry for such spaces. There exist only three types of homogeneous and isotropic spaces with simple topology: (a) flat space, (b) a three-dimensional sphere of constant positive curvature, and (c) a three-dimensional hyperbolic space of constant negative curvature [7].

Many interesting results on curves of AW(k)-type have been obtained by many mathematicians [1,3,4,5,6]. Also, Bertrand curves have been studied in [8] and [11].

In this paper, we have done a study about some special curves in Isotropic Space I_3^1 . However, to the best of author's knowledge, Bertrand and Mannheim curves of AW(k)-type has not been presented in Isotropic Space I_3^1 . Thus, the study is proposed to serve such a need.

Our paper is organized as follows. In section 2, the basic notions and properties of a Frenet curve are reviewed. In section 3, we study curves of AW(k)-type in Isotropic Space I_3^1 . We also study Bertrand and Mannheim curves of AW(k)-type in section 4.

2. BASIC NOTIONS AND PROPERTIES

Let $\alpha : I \rightarrow I_3^1, I \subset IR$ be a curve given by

$$\alpha(t) = (x(t), y(t), z(t)),$$

where $x, y, z \in C^3$ (the set of three times continuously differentiable functions) and t run through a real interval [9].

Let α be a curve in I_3^1 , parameterized by arc length $t = s$, given in coordinate form

$$\alpha(s) = (s, y(s), z(s)). \tag{1}$$

In [9], the curvature $\kappa(s)$ and the torsion $\tau(s)$ are defined by

$$\kappa(s) = x' y'' - y' x''$$

$$\tau(s) = \frac{\det(\alpha'(s), \alpha''(s), \alpha'''(s))}{\kappa^2(s)} \tag{2}$$

and associated moving trihedron is given by

$$t(s) = \alpha'(s)$$

$$n(s) = \frac{1}{\kappa(s)} \alpha''(s) \tag{3}$$

$$b(s) = (0, 0, 1)$$

The vectors t, n, b are called tangent vector field, principal normal vector field and binormal vector field of the curve α , respectively. For their derivatives the following Frenet formulas hold

$$\begin{aligned} t'(s) &= \kappa(s)n(s) \\ n'(s) &= -\kappa(s)t(s) + \tau(s)b(s) \\ b'(s) &= 0 \end{aligned} \tag{4}$$

Scalar product in the Isotropic space I_3^1 is defined by

$$\langle X, Y \rangle = x_1y_1 + x_2y_2 \tag{5}$$

where $X = (x_1, x_2, x_3)$ and $Y = (y_1, y_2, y_3)$.

If $x_1y_1 + x_2y_2 = 0$, then $\langle X, Y \rangle = x_3y_3$.

The isotropic norm of a vector $X = (x_1, x_2, x_3)$ is defined by

$$\|X\| = \sqrt{x_1^2 - x_2^2}$$

where \sim on the vector denotes the canonical projection of the vector to the base plane $x_3 = 0$. If $\|X\| = 0$, i.e. if X is an isotropic vector, then the supplementary invariant called range of the vector X is introduced

$$[X] = x_3.$$

If $\|X\| \neq 0$, then X called Euclidean vector [10].

From now on in calculations, " \tilde{X} " canonical projection of the vectors are denoted as " X ".

According to [1], one can calculate the followings:

Proposition 2.1 Let α be a Frenet curve of I_3^1 of osculating order 3 then we have

$$\begin{aligned} \alpha'(s) &= t(s) \\ \alpha''(s) &= t'(s) = \kappa(s)n(s) \end{aligned} \tag{6}$$

$$\alpha'''(s) = -\kappa^2(s)t(s) + \kappa'(s)n(s) + \kappa(s)\tau(s)b(s) \tag{7}$$

$$\begin{aligned} \alpha^{IV}(s) &= -3\kappa(s)\kappa'(s)t(s) + [\kappa''(s) - \kappa^3(s)]n(s) \\ &\quad + [2\kappa'(s)\tau(s) + \kappa(s)\tau'(s)]b(s) \end{aligned} \tag{8}$$

Notation. Let us write

$$N_1(s) = \kappa(s)n(s) \tag{9}$$

$$N_2(s) = \kappa'(s)n(s) + \kappa(s)\tau(s)b(s) \tag{10}$$

$$N_3(s) = [\kappa''(s) - \kappa^3(s)]n(s) + [2\kappa'(s)\tau(s) + \kappa(s)\tau'(s)]b(s) \tag{11}$$

Corollary 2.2 $\alpha'(s), \alpha''(s), \alpha'''(s)$ and $\alpha^{IV}(s)$ are linearly dependent if and only if $N_1(s), N_2(s)$ and $N_3(s)$ are linearly dependent.

Theorem 2.3 Let α be a Frenet curve of I_3^1 of osculating order 3 then

$$N_3(s) = \langle N_3(s), N_1^*(s) \rangle N_1^*(s) + \langle N_3(s), N_2^*(s) \rangle N_2^*(s) \tag{12}$$

where

$$N_1^*(s) = \frac{N_1(s)}{\|N_1(s)\|}, \quad N_2^*(s) = \frac{N_2(s) - \langle N_2(s), N_1^*(s) \rangle N_1^*(s)}{\|N_2(s) - \langle N_2(s), N_1^*(s) \rangle N_1^*(s)\|}. \tag{13}$$

3. CURVES OF AW(K)-TYPE

Definition 3.1 (see , cf.[1]) Frenet curves (of osculating order 3) are
i) of type weak AW(2) if they satisfy

$$N_3(s) = \langle N_3(s), N_2^*(s) \rangle N_2^*(s), \tag{14}$$

ii) of type weak AW(3) if they satisfy

$$N_3(s) = \langle N_3(s), N_1^*(s) \rangle N_1^*(s). \tag{15}$$

Proposition 3.2 Let \mathcal{C} be a Frenet curve of order 3. If \mathcal{C} is of type weak AW(2) then

$$\kappa''(s) - \kappa^3(s) = 0. \tag{16}$$

Corollary 3.3 Let \mathcal{C} be a Frenet curve of type weak AW(2). If \mathcal{C} is a plane curve then

$$\kappa(s) = \frac{\sqrt{2}}{s+c}; \quad c = const. \tag{17}$$

Proposition 3.4 Let \mathcal{C} be a Frenet curve of order 3. If \mathcal{C} is of type weak AW(3) then

$$2\kappa'(s)\tau(s) + \kappa(s)\tau'(s) = 0. \tag{18}$$

Corollary 3.5 Let \mathcal{C} be a Frenet curve of type weak AW(3). Then

$$\tau(s) = \frac{c}{\kappa^2(s)}; \quad c = const. \tag{19}$$

Definition 3.6 (see , cf.[1]) Frenet curves are

i) of type AW(1) if they satisfy

$$N_3(s) = 0, \tag{20}$$

ii) of type AW(2) if they satisfy

$$\|N_2(s)\|^2 N_3(s) = \langle N_3(s), N_2(s) \rangle N_2(s), \tag{21}$$

iii) of type AW(3) if they satisfy

$$\|N_1(s)\|^2 N_3(s) = \langle N_3(s), N_1(s) \rangle N_1(s). \tag{22}$$

Theorem 3.7 Let \mathcal{C} be a Frenet curve of order 3. Then \mathcal{C} is of type AW(1) if and only if

$$\kappa''(s) - \kappa^3(s) = 0 \tag{23}$$

and

$$\tau(s) = \frac{c}{\kappa^2(s)}; \quad c = \text{const.} \tag{24}$$

Proof. Let α be a curve of type AW(1). Then from (11) and (20) we have

$$[\kappa''(s) - \kappa^3(s)]n(s) + [2\kappa'(s)\tau(s) + \kappa(s)\tau'(s)]b(s) = 0.$$

Furthermore, since $n(s)$ and $b(s)$ are linearly independent, we get (23) and (24).

The converse statement is trivial. Hence our theorem is proved.

Corollary 3.8 Every plane curve of type AW(1) is also of type weak AW(2).

Theorem 3.9 Let \mathcal{C} be a Frenet curve of order 3. Then α is of type AW(2) if and only if

$$2[\kappa'(s)]^2 \kappa(s)\tau^2(s) + \kappa^2(s)\tau(s)\kappa'(s)\tau'(s) + \kappa^5(s)\tau^2(s) - \kappa''(s)\kappa^2(s)\tau^2(s) = 0 \tag{25}$$

and

$$2[\kappa'(s)]^3 \tau(s) + [\kappa'(s)]^2 \kappa(s)\tau'(s) + \kappa^4(s)\kappa'(s)\tau(s) - \kappa'(s)\kappa''(s)\kappa(s)\tau(s) = 0 \tag{26}$$

Proof. If α curve is of type AW(2), (21) holds on α . Substituting (10) and (11) into (21), we have (25) and (26).

Theorem 3.10 Let α be a Frenet curve of order 3. Then α is of type AW(3) if and only if

$$2\kappa^2(s)\kappa'(s)\tau(s) + \kappa^3(s)\tau'(s) = 0. \tag{27}$$

Proof. Since α is of type AW(3), (22) holds on α . So substituting (9) and (11) into (22), we have (27).

4. BERTRAND CURVES AND MANNHEIM CURVES OF AW(K)-TYPE

In this section, we give the characterizations of Bertrand and Mannheim Curves of AW(k)-type.

Remark 4.1 Let α be a Frenet curve of order 3 of I_3^1 . For $\tau(s) \neq 0$, α is a Bertrand curve if and only if there exist a linear relation

$$A\kappa(s) + B\tau(s) = 1 \tag{28}$$

where A, B are non-zero constant and $\kappa(s)$ and $\tau(s)$ are the curvature functions of α [9].

Corollary 4.2 Suppose that $\kappa(s) \neq 0$ and $\tau(s) \neq 0$. Then α is a Bertrand curve if and only if there exist a non-zero real number A such that [2]

$$A[\tau'(s)\kappa(s) - \kappa'(s)\tau(s)] - \tau'(s) = 0. \tag{29}$$

Theorem 4.3 Let $\alpha : I \rightarrow I_3^1$ be a Bertrand curve with $\kappa(s) \neq 0$ and $\tau(s) \neq 0$. Then α is of type AW(2) if and only if there is a non-zero real number A such that

$$2[\kappa'(s)]^2 \kappa(s)\tau^2(s) + A\kappa^3(s)\kappa'(s)\tau(s)\tau'(s) - \kappa^2(s)[\kappa'(s)]^2 \tau^2(s) + \kappa^5(s)\tau^2(s) - \kappa''(s)\kappa^2(s)\tau^2(s) = 0 \tag{30}$$

and

$$2[\kappa'(s)]^3 \tau(s) + A\kappa^2(s)[\kappa'(s)]^2 \tau'(s) - \kappa(s)[\kappa'(s)]^3 \tau(s) + \kappa^4(s)\kappa'(s)\tau(s) - \kappa'(s)\kappa''(s)\kappa(s)\tau(s) = 0 \tag{31}$$

Proof. Since α is of type AW(2), (25) and (26) holds and since α is a Bertrand curve, (29) equality holds. If both of these equations are considered, (30) and (31) are obtained.

Theorem 4.4 Let $\alpha : I \rightarrow I_3^1$ be a Bertrand curve with $\kappa(s) \neq 0$ and $\tau(s) \neq 0$. Then α is of type AW(3) if and only if

$$2\kappa^2(s)\kappa'(s)\tau(s) + A\kappa^4(s)\tau'(s) - \kappa^3(s)\kappa'(s)\tau(s) = 0. \tag{32}$$

Proof. Now suppose that $\alpha : I \rightarrow I_3^1$ be a Bertrand curve of type AW(3) with $\kappa(s) \neq 0$ and $\tau(s) \neq 0$. Then the equation (27) and (29) hold on α . Thus, we get (32).

Definition 4.5 Let α be a curve in I_3^1 . If its principal normal vector field is the binormal vector field of another curve, then the curve α is called Mannheim curve in I_3^1 .

Theorem 4.6 Let α be a curve in I_3^1 . Then α is Mannheim curve if and only if its curvature

$$\kappa(s) = c; \quad c = const. \tag{33}$$

Proof. Let $\alpha = \alpha(s)$ be a Mannheim curve in I_3^1 . Let us denote of Frenet Frame of the curve α by $\{t_\alpha(s), n_\alpha(s), b_\alpha(s)\}$. The curve $\bar{\alpha}(s)$ is parametrized by arclength s as

$$\bar{\alpha}(s) = \alpha(s) + c_1(s)n(s) \tag{34}$$

for some functions $c_1(s) \neq 0$. Differentiating (34) with respect to s , we find

$$\overline{\alpha}'(s) = (1 - c_1(s)\kappa(s))t(s) + c_1'(s)n(s) + c_1(s)\tau(s)b(s). \tag{35}$$

Since the binormal vector of $\overline{\alpha}(s)$ is linearly dependent with principal normal vector of $\alpha(s)$, we have

$$c_1'(s) = 0.$$

Hence $c_1(s) = \text{const.}$ The second derivative $\overline{\alpha}''(s)$ with respect to s is

$$\overline{\alpha}''(s) = -c_1(s)\kappa'(s)t(s) + [\kappa(s) - c_1(s)\kappa^2(s)]n(s) + c_1(s)\tau'(s)b(s). \tag{36}$$

Since $n(s)$ is the binormal direction of $\overline{\alpha}(s)$, we have

$$\kappa(s) - c_1(s)\kappa^2(s) = 0. \tag{37}$$

From (37), we get

$$\kappa(s) = c \tag{38}$$

where $c = \frac{1}{c_1(s)}$.

Conversely, let $\overline{\alpha}(s)$ be a curve in I_3^1 with $\kappa(s) = \frac{1}{c_1(s)}$. Then the curve

$$\overline{\alpha}(s) = \alpha(s) + c_1(s)n(s)$$

has binormal direction $n(s)$. It follows that $\alpha(s)$ is a Mannheim curve which proves the theorem.

Theorem 4.7 Let α be a Mannheim curve in I_3^1 . Then \mathcal{C} is of type AW(1) if and only if

$$\tau(s) = \text{const.} \tag{39}$$

Proof. Considering Theorem 4.6. in Theorem 3.7., we get (39). Hence the proof is completed.

Theorem 4.8 Let α be a Mannheim curve in I_3^1 . Then α is of type AW(2) if and only if

$$\tau(s) = 0. \tag{40}$$

Proof. Considering Theorem 4.6. in Theorem 3.9., we get (40). Hence our theorem is proved.

Theorem 4.9 Let α be a Mannheim curve in I_3^1 . Then α is of type AW(3) if and only if

$$\tau(s) = \text{const.} \tag{41}$$

Proof. Considering Theorem 4.6. in Theorem 3.10., we get (41). Hence the proof is completed.

Example 4.10 Let α be a curve in I_3^1 given by

$$\alpha(u) = \left(a \cos \frac{u}{a}, a \sin \frac{u}{a}, 0 \right)$$

Then we have

$$\alpha'(u) = \left(-\sin \frac{u}{a}, \cos \frac{u}{a}, 0 \right)$$

$$\alpha''(u) = \left(-\frac{1}{a} \cos \frac{u}{a}, -\frac{1}{a} \sin \frac{u}{a}, 0 \right)$$

Using (2) equality, we get $\kappa(s) = \frac{1}{a}$, $\tau(s) = 0$. $\kappa(s)$ and $\tau(s)$ hold on Theorems of 3.9, 3.10, 4.3, 4.4 and 4.8.

5. CONCLUSION

It is well-known that isotropic spaces are very important in physics and mathematics. Because isotropic spaces have the largest possible symmetry group: in three dimensions there are three independent translations and three rotations.

In this study, AW(k)-type curves are examined in Isotropic space I_1^3 .

It is hoped that this study serves researchers who carry out research especially in geometry and mathematical physics.

ACKNOWLEDGEMENTS

We would like to thank for constructive criticisms provided by Editorial Board and anonymous reviewers whose insights improved the quality of the article.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Arslan K, Özgür C. Curves and surfaces of AW(k)-type. Geometry and topology of submanifolds. IX (Valenciennes/Lyan/Leuven, 1997), 21--26. River Edge, [NJ]: World Scientific Publishing; 1999.
2. Izumiya S, Takeuchi N. New special curves and developable surfaces. Turkish J. Math. 2004;28(2):153-163.
3. Külahcı M, Bektaş M, Ergüt M. Curves of AW(k)-type in three-dimensional null cone. Phys Lett A. 2007;371:275--7.
4. Külahcı M, Bektaş M, Ergüt M. On harmonic curvatures of null curves of AW(k)-type in Lorentzian space. Z Naturforsch. 2008;63a:248-52.
5. Külahcı M, Bektaş M, Ergüt M. On harmonic curvatures of a Frenet curve in Lorentzian space. Chaos, Solitons and Fractals. 2009;41:1668-1675.

6. K lahcı M, Erg t M. Bertrand curves of AW(k)-type in Lorentzian space. *Nonlinear Anal.* 2009;70(4):1725-1731.
7. Mukhanov VF. *Physical foundations of cosmology*. Cambridge University Press, New York; 2005.
8.  grenmiŐ AO,  ztekin H, Erg t M. Bertrand Curves in Galilean Space and their Characterizations. *Kragujevac J. Math.* 2009;(32):139-147.
9. Sachs H. *Isotrope Geometrie des Raumes*. Braunschweig: Vieweg; 1990.
10. Sipus ZM, Divjak B. Mappings of Ruled Surfaces in Simply Isotropic Space I_3^1 that preserve the generators. *Monatsh. Math.* 2003;139:235-245.
11. Yıldırım Yılmaz M, BektaŐ M. General properties of Bertrand curves in Riemann-Otsuki space. *Nonlinear Anal.* 2008;69(10):3225-3231.

  2013 Ogrenmis et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history.php?iid=224&id=4&aid=1451>